

---

# NASA/ADS

## The Formation of Magnetic Decreases $MDs$ in Interplanetary Space: Ulysses

Show affiliations

**Tsurutani, B.; Guarnieri, F.; Echer, E.**

Magnetic Decreases are decreases in the magnitude of the ambient magnetic field. The displaced plasma pressure is supplanted by anisotropic  $T_{perp}/T_{par} > 1$ , locally heated plasma so that there is pressure balance across the MD structures. Anisotropic spatial distributions of MDs are identified to further understand the conditions under which MDs form/evolve. We will show that Ulysses fast-latitude scans through corotating solar wind streams are good places to search for such spatial anisotropies. The properties of MDs within CIRs and high speed streams will be reported. A general generation mechanism will be described.

**Publication:**

American Geophysical Union, Fall Meeting 2007, abstract id.SH31B-04

**Pub Date:**

December 2007

**Bibcode:**

2007AGUFMSH31B..04T

**Keywords:**

2102 Corotating streams; 2134 Interplanetary magnetic fields;  
2159 Plasma waves and turbulence

 Feedback/Corrections? ([http://adsabs.harvard.edu/adsfeedback/submit\\_abstract.php?bibcode=2007AGUFMSH31B..04T](http://adsabs.harvard.edu/adsfeedback/submit_abstract.php?bibcode=2007AGUFMSH31B..04T))