

Integrated Nitrogen Catchment model (INCA) applied to a tropical catchment in the Atlantic Forest, São Paulo, Brazil

M. Ranzini^{1,2}, M.C. Forti^{2,3}, P.G. Whitehead⁴, F.C.S. Arcova¹, V. de Cicco¹ and A.J. Wade⁴

¹Instituto Florestal, P.O.Box 1322, 01059-970, São Paulo, Brazil

²Instituto Nacional de Pesquisas Espaciais INPE, P.O.Box 515, 12245-970, S.J.dos Campos, Brazil

³Núcleo de Pesquisa em Geoquímica e Geofísica da Litosfera, University of São Paulo, P.O. Box 9, 13418-900, Piracicaba, Brazil

⁴Aquatic Environments Research Centre, School of Human and Environmental Sciences, University of Reading, Reading, RG6 6AB, UK

Email for corresponding author: ranzini@ig.com.br

Abstract

Stream-water flows and in-stream nitrate and ammonium concentrations in a small (36.7 ha) Atlantic Forest catchment were simulated using the Integrated Nitrogen in Catchments (INCA) model version 1.9.4. The catchment, at Cunha, is in the Serra do Mar State Park, SE Brazil and is nearly pristine because the nearest major conurbations, São Paulo and Rio, are some 450 km distant. However, intensive farming may increase nitrogen (N) deposition and there are growing pressures for urbanisation. The mean-monthly discharges and NO₃-N concentration dynamics were simulated adequately for the calibration and validation periods with (simulated) loss rates of 6.55 kg.ha⁻¹ yr⁻¹ for NO₃-N and 3.85 kg.ha⁻¹ yr⁻¹ for NH₄-N. To investigate the effects of elevated levels of N deposition in the future, various scenarios for atmospheric deposition were simulated; the highest value corresponded to that in a highly polluted area of Atlantic Forest in São Paulo City. It was found that doubling the atmospheric deposition generated a 25% increase in the N leaching rate, while at levels approaching the highly polluted São Paulo deposition rate, five times higher than the current rate, leaching increased by 240%, which would create highly eutrophic conditions, detrimental to downstream water quality. The results indicate that the INCA model can be useful for estimating N concentration and fluxes for different atmospheric deposition rates and hydrological conditions.

Keywords: Atlantic Forest, tropics, N-modelling, hydrology, nitrogen dynamics, INCA, atmospheric deposition, Cunha

Introduction

The Atlantic Forest (Mata Atlântica) is a montane rainforest along the Brazilian Atlantic Coast. The forest lies parallel to the eastern coast of Brazil from 25°S (Santa Catarina state) to 5°S (Rio Grande do Norte state), with an altitude range from 800–1500 m. Formerly the Mata Atlântica occupied 15% of the country, but the biome has suffered from human modifications, beginning in the 16th Century (Fundação SOS Mata Atlântica, 2005). Figure 1a and b shows the original and the present territorial distributions of the Atlantic Forest. The rainforest is characterised by a hot, humid climate with dense and diverse vegetation. Following the colonisation of Brazil, less than 4% of the original total area of the Atlantic Forest now remains because of wood extraction and urbanisation, including the Metropolitan Area of São Paulo (19 million inhabitants) (Emplasa, 2005). In

São Paulo State, the remnant is only 15% of the original area (Fundação SOS Mata Atlântica, 2005).

The desire to conserve and manage the remainder of the Atlantic Forest in a sustainable way requires a better understanding of the hydrogeochemical fluxes, particularly those for nitrogen (N) since it is a key nutrient. Moreover, given the likely continuation of urbanisation and agricultural expansion in the region, the effects of increased inputs of pollutants from atmospheric deposition on the N budget must be quantified. The N cycle is complex and its processes are not well known in tropical areas. Although N is not limiting for vegetation nutrition in this region, it is important to control the trophic level of water bodies (Neal, 2002); managing N inputs to catchments is, therefore, a key requirement. Concentrations of nitrate in rivers reflect the integration of diverse sources within the catchment above




Fig. 1. The Brazilian Atlantic Forest, showing original distribution (left) and present distribution (right). (Fundação SOS Mata Atlântica, 2005).

the background inputs (the mineralisation and nitrification of organic N in the soils). These additional sources include fertiliser inputs, atmospheric deposition and sewage. The fluxes of nitrogen through catchments will be influenced by climatic and seasonal factors (e.g. drought and temperature), and by land use and management practices such as afforestation, clear-felling, liming, ploughing and grazing. The modifications that result from anthropogenic activity can interfere with hydrogeochemical processes and, potentially, threaten water resources.

The complexity of the processes involved requires models to predict how atmospheric deposition, soil use, ecosystem management and climate change all affect N concentrations and loads in rivers. In particular, with representative field measurements, models can help understanding of how nitrogen is cycled and transported and how the key factors and processes that control its dynamics can be determined. Moreover, for environmental management, models are needed to provide quantitative estimates of how a system will respond to changes in pollutant inputs.

A new version of the process-based INCA model (Whitehead *et al.*, 1998a, b; Wade *et al.*, 2002) was chosen for this study because it accounts for the fluxes and storages of nitrogen in both land and stream components of a system, tracking the inputs from the atmosphere and from fertiliser through catchment soils to the river. It also simulates the effects of spatial variations in the land use and hydrology of a river system. The INCA model has been tested in river systems throughout Europe and while it has been found to be applicable across a wide range of spatial and temporal scales (Neal, 2002; Wade *et al.*, 2002), it has yet to be applied in tropical or subtropical regions, so the present study pioneers the application of an established model of river-system N dynamics to a different ecosystem. The objectives of this study are (1) to use the model as a learning tool to understand the hydro-geochemical processes controlling the nitrogen dynamics of a small tropical forest catchment and (2) to use the calibrated model to quantify the likely response of the catchment to predicted changes in N deposition, a potential threat to water quality in the Atlantic Forest region.

Material and methods

The small (36.7 ha) Cunha catchment, in the Serra do Mar State Park, is some 15 km from the Atlantic Ocean in south-east Brazil, (lat. 23°13' S, long. 45°02' W), and lies at elevations between 1000–1200 m.a.s.l. (Fig. 2). The mean annual air temperature is 16.5°C and the mean annual total precipitation is 2205 mm (Cicco, 2004). The bedrock is gneiss and crystalline schist of Precambrian age. Oxysols (FAO-System) comprise the dominant soil coverage, with Gleyic Cambisols close to the creeks (Furian and Pfeifer, 1986).

The INCA model is a process-based model of the nitrogen cycle in plant/soil and in-stream systems. The study catchment was considered as a single reach. The driving data for the INCA model, soil moisture deficit, hydrologically effective rainfall (the fraction of precipitation that contributes directly to the runoff) and air temperature were compiled from April 01 2000 to March 31 2004 and daily potential evapotranspiration was estimated using Camargo's method (1971). Meteorological measurements

of air temperature, air humidity, solar radiation and wind speed have been made since January 1998 at a weather station 2 km from the catchment while, since October 1986, daily precipitation has been measured with tipping-bucket rain-gauges installed in three clearings. The hydrologically effective rainfall was calculated as the real precipitation minus the evapotranspiration, taking into account the water retention by the soils (Fujieda *et al.*, 1997; Ranzini, 2002).

Since October 1986, discharge from the catchment has been measured with a 90° sharp crested weir with a stilling pond and a continuous record of stage is available. The rating curve for the weir determined by current metering under contrasting flow conditions (Arcova, 1996). Weekly stream water samples were taken from May 2000 to February 2004 at the outlet of the catchment. Bulk and wet-only rainfall water samples were collected during the same period. However, after rainfall collectors had been exposed for 15 days, if insufficient water had accumulated, the collection funnel and bottles were replaced by clean ones to minimise the influence of dry deposition on the sample. A wet-only collector (MTX-Italy) was deployed simultaneously with the bulk collectors and the two sets of measurements were used to estimate the effect of dry deposition on the rainfall chemistry (Forti *et al.*, 2005).

The parameters used in the INCA model are shown in Table 1. Because the forest under study is natural and mature, the parameters related to fertilisation are set to zero in the model. The maximum soil moisture deficit, the initial soil water flow and the drainage volume were determined empirically from another catchment (D) located in the same area (Ranzini *et al.*, 2004). Other parameters associated with soil reactive zone, groundwater zone, initial surface flow and initial sub-surface flow, and the base flow index, were either calculated from analyses of the continuous stage record or measured directly. Parameters relating to the velocity flow relationship in the stream were adjusted during the calibration process.

The initial values of the parameters relating to denitrification, nitrogen fixation, nitrification, N-mineralisation, immobilisation, and in-stream denitrification and nitrification rates were obtained from an application of INCA to forested catchments in Germany (Langusch and Matzner, 2002). These values were used because there was no comparable information for the present study area. After an initial calibration procedure, the parameters were adjusted to fit the observed nitrogen dynamics as closely as possible. The dry and wet deposition rates for nitrate and ammonium were obtained from observations (Forti *et al.*, 2005).




Fig. 2. Location of the Cunha Forest Hydrological Laboratory.

Table 1. Parameters of the INCA model used for the modelling.

Parameters obtained for INCA V1.9.4 calibration-CUNHA

<u>Land Phase</u>		<u>Sub-Model</u>	
Volume Constants		Soil temperature	
Ratio of total to available water in soil	2 Φ	Difference between maximums	6°C
Maximum Groundwater effective depth	60 m	Thermal conductivity in soil	0.7W.m ² .°C
Proportion of filed spaces	0.5 Φ	Specific heat capacity due to freezer/thous	6.6 J.m ³ .°C
		Snow pack effect – disabled	
		Temperature response – disabled	
		Snow pack - disabled	
		Fertilizer	
		Addition – zero	
Time constants		Plant growth	
Direct runoff residence time	0.5 d	Growth season start day	1 Julian day
Soil water residence time	3 d	Growth period	730 d
Ground water residence time	150 d	Nitrate uptake rate	0.01 m.d ⁻¹
		Ammonium uptake rate	0 m.d ⁻¹
		Max N uptake	100 kgN.ha ⁻¹ .yr ⁻¹
		Growth curve offset	0.66 Φ
		Growth curve amplitude	0.34 Φ
Processes – Soil water		<u>Instream parameters</u>	
Denitrification rates	0.01 m.d ⁻¹	Initial values	
Fixation rates	0 kgNha ⁻¹ .d ⁻¹	Flow	0.0464 m ³ .s ⁻¹
Nitrification rates	0.1m.d ⁻¹	Nitrate	0.3 mgN.L ⁻¹
Mineralisation rates	0.15 kgNha ⁻¹ .d ⁻¹	Ammonium	0.2 mgN.L ⁻¹
Immobilisation rates	0 m.d ⁻¹	Minimum temperature	0°C
		<u>Reach parameters</u>	
Processes - Ground water		Length	
Denitrification rates	0 m.d ⁻¹	a	920 m
		b	0.04 m ²
		Denitrification	0.67 Φ
		Nitrification	0.05.d ⁻¹
Initial Values – Direct runoff		<u>Sub catchment Parameters</u>	
Flow	0.001 m ³ .s ⁻¹	Physical attributes	
Nitrate	0.3 mgN.L ⁻¹	Area	0.367 km ²
Ammonium	2.9 mgN.L ⁻¹	Base flow	0.52 Φ
Initial values – Soil water		Direct runoff – saturation excess control	
Flow	0.1 m ³ .s ⁻¹	Threshold soil zone flow	0.06 m ³ .s ⁻¹
Nitrate	2.0 mgN.L ⁻¹	Infiltration excess control	
Ammonium	1.98 mgN.L ⁻¹	Rainfall excess propostion	0.05 Φ
Initial values – Ground water		Max infiltration rate	5 mm.d ⁻¹
Flow	0.05 m ³ .s ⁻¹	Deposition	
Nitrate	0.1 mgN.L ⁻¹	Nitrate dry	0.01 kgN.ha ⁻¹ .yr ⁻¹
Ammonium	0.3 mgN.L ⁻¹	Nitrate wet	9.0 kgN.ha ⁻¹ .yr ⁻¹
		Ammonium dry	0.0 kgN.ha ⁻¹ .yr ⁻¹
		Ammonium wet	8.0 kgN.ha ⁻¹ .yr ⁻¹
Threshold – soil moisture deficit		Land use	
Maximum soil moisture deficit	10 mm	Forest 100%	
Threshold – sustainable flow		Calibration period	
Minimum soil water flow	9999 m ³ .L ⁻¹	1/4//2000 – 31/3/2002	
Minimum ground water flow	9999 m ³ .L ⁻¹	Validation period	
		1/4/2002 – 27/2/2004	
Threshold – processes temperature			
Not activated			

Model calibration

Four years of discharge and NO₃-N concentration measurements were available and a split sample test used data for April 2000 to March 2002 for calibration and for April 2002 to March 2004 for validation of the model. The model calibration followed three steps following Jarvie *et al.* (2002):

1. *Hydrology.* To simulate the correct dilution of the nitrogen mass in the stream and the timing of the inputs from the land to the stream, it is important to simulate the hydrology accurately. The water residence time constants in the unsaturated and saturated zones were taken from the work of Fujieda *et al.* (1997).
2. *Initial conditions.* The second step in the calibration procedure was the adjustment of the NO₃-N and NH₄-N initial concentrations in the soil, groundwater and in-stream components, so that the initial flow NO₃-N concentrations matched observed in-stream concentrations.
3. *Processes rates.* Parameters relating to soil nitrogen processes (rates of NH₄-N immobilisation, NH₄-N nitrification, NO₃-N denitrification, NH₄-N mineralisation, and plant NH₄-N and NO₃-N uptake) as well as in-stream rates of denitrification and NH₄-N nitrification were adjusted so that the simulated N annual fluxes and leaching and within-catchment processes had

values comparable with those in the literature, and the simulated daily NO₃-N concentrations matched those actually observed.

The calibrated parameters, shown in Table 1, are consistent with those found in other studies with the INCA model.

Results

CALIBRATION

The observed and simulated monthly mean discharges (April 2000 to March 2004) were similar (Fig. 3a, b): the simulated mean value of 82 mm approximated the observed mean of 87 mm, the coefficient of determination of the correlation was $r^2 = 0.74$ and the efficiency (Nash and Sutcliffe, 1970) was 0.72.

Rainfall in the region is seasonal: a relatively dry period with low temperatures from June to August corresponds to the winter months and high rainfall and higher temperatures between December and March correspond to the summer season. For most of the year, the soil moisture deficit is less than 4 mm; in an adjacent catchment, after a period of 20 days without rain, the soil moisture profile was approximately the field capacity of the soil (Ranzini, 2002). The seasonal soil moisture deficit pattern differed markedly from the pronounced seasonal differences found in forested catchments in temperate regions modelled previously with




Fig. 3a. Observed and simulated mean monthly flow ($m^3 \cdot s^{-1}$), and nitrate and ammonium stream water concentrations ($mg \cdot L^{-1}$) for the calibration period (April 2000 to March 2002).




Fig. 3b. Observed and simulated mean monthly flow ($m^3 \cdot s^{-1}$), and nitrate and ammonium stream water concentrations ($mg \cdot L^{-1}$) for the validation period (April 2002 to March 2004).

INCA (Langush and Matzner, 2002; Bastrup-Birk and Gundersen, 2004). The pattern in temperate regions results from the combination of high temperatures and low rainfall during the summer (Whitehead *et al.*, 1998b, 2002).

The dynamics of the in-stream concentrations of NO_3 -N and NH_4 -N were well explained by the model (Fig. 3a,b). Mean concentrations of NO_3 -N and NH_4 -N, $0.43 mg L^{-1}$ and $0.23 mg L^{-1}$ respectively were consistent with modelled mean values of $0.50 mg L^{-1}$ and $0.24 mg L^{-1}$ respectively.

Leaching of NH_4 -N, overestimated with the present parameterisation (Table 2), may be attributed to a sampling bias as most values were determined during periods of base-flow. Nevertheless, the NH_4 -N and NO_3 -N concentrations are normal for a forested catchment (Langusch and Matzner, 2002) and are consistent with values of $0.14 mg L^{-1}$ and $0.66 mg l^{-1}$ for NH_4 -N and NO_3 -N, respectively, found in the Amazon basin by Forti *et al.* (2000).

The NO_3 -N and NH_4 -N modelled leaching rates of $6.5 kg N ha^{-1} yr^{-1}$ and $3.85 kg ha^{-1} yr^{-1}$ respectively, are 20% and 40% higher than those observed (Table 2). These differences may be due to the sampling procedure as well as to a lack of good data for the nitrogen processes (Fig. 3a, b). Mineralisation and nitrification rates obtained from the modelling, $10.5 kg ha^{-1} yr^{-1}$ and $15.8 kg ha^{-1} yr^{-1}$, are lower than those observed for a soil chronosequence (assuming a mean density of $1.3 g cm^{-3}$ and 1.8 m depth) for the Amazon

(Neill *et al.*, 1997) where values of mineralisation were from $17.9 kg ha^{-1} yr^{-1}$ to $53.3 kg ha^{-1} yr^{-1}$ and of nitrification from 20.2 and $55.1 kg ha^{-1} yr^{-1}$.

MODEL VALIDATION

To validate the model, the parameters determined in the INCA calibration for the period from April 2000 to March 2002 were used to simulate results for the period from April 2002 to March 2004. In this way, the modelled hydrology and NO_3 -N and NH_4 -N fluxes and concentrations could be tested using a different data set by comparing the simulated and observed values of the daily discharge and daily NO_3 -N and NH_4 -N concentrations for the validation period. There was a good match between the simulated and observed discharge, as well as of nitrate and ammonium dynamics (Fig. 3a,b). As in the calibration mode, in the rainy period when peak discharges occur, NO_3 -N and NH_4 -N concentrations also peak, thus reproducing the observed seasonality.

ENVIRONMENTAL CHANGES SIMULATION WITH DIFFERENT ATMOSPHERIC DEPOSITION SCENARIOS

Although the catchment under study lies within a

Table 2. Annual load data for the different parameters for the observed data and the INCA simulations.

Land Use Group-Forest	Observed	Calibration	Validation	Scenario 1	Scenario 2	Scenario 3	Scenario 4	Scenario 5
Nitrate Wet deposition kgN.ha ⁻¹ .yr ⁻¹		9	9	18	27	36	45	54
Reactive Zone Flow m ³ .km ²		1981	15300	1981	1981	1981	1981	1981
Groundwater flow m ³ .km ²		2.96E+07	2.96E+07	2.96E+07	2.96E+072.96E+072.96E+072.96E+07	2.96E+072.96E+072.96E+072.96E+07	E+07	2.96E+07
Nitrate-N Total Load kgN.ha ⁻¹ .yr ⁻¹	10.8	9.09	9.01					
Ammonium-N Total Load kgN.ha ⁻¹ .yr ⁻¹	9.3	8.07	8.0	18.15	27.23	36.31	45.38	54.45
Nitrate-N Leaching kgN.ha ⁻¹ .yr ⁻¹	5.22	6.5	6.8	8.26	9.83	11.41	12.99	14.57
Ammonium-N Leaching kgN.ha ⁻¹ .yr ⁻¹	2.38	3.85	3.97	5.19	6.39	7.60	8.80	10.01
Nitrate-N Uptake kgN.ha ⁻¹ .yr ⁻¹		6.24	6.39	8.04	9.66	11.28	12.91	14.53
Ammonium-N Uptake kgN.ha ⁻¹ .yr ⁻¹		0	0	0	0	0	0	0
Ammonium-N Mineralization kgN.ha ⁻¹ .yr ⁻¹		10.5	11.15	10.5	10.5	10.5	10.5	10.5
Ammonium-N Nitrification kgN.ha ⁻¹ .yr ⁻¹		15.79	16.24	23.69	30.8	37.91	45.02	52.15
Nitrate-N Denitrification kgN.ha ⁻¹ .yr ⁻¹		9.92	10.09	12.91	15.61	18.3	21.0	23.70
Nitrate-N Fixation kgN.ha ⁻¹ .yr ⁻¹		0	0	0	0	0	0	0
Ammonium-N Retention kgN.ha ⁻¹ .yr ⁻¹		0	0	0	0	0	0	0

conservation unit (State Park of Serra do Mar), it is potentially vulnerable to changes in N emissions from the surrounding regions, particularly from intensive cattle ranching and increasing industrial development in the Paraíba do Sul river valley, with a corresponding increase in human population. To simulate these changes, a range of scenarios was chosen: 100 to 500% increase in total N deposition above present baseline levels for Cunha. The highest values correspond to the deposition in a highly impacted part of the Atlantic Forest: the Fontes do Ipiranga State Park – PEFI, a forested area of 526.4 ha surrounded by the metropolitan region of São Paulo with a mean N deposition rate of 50.8 kg ha⁻¹ yr⁻¹ (Forti *et al.*, 2005), around six times the NO₃-N wet deposition value used for calibration.

Table 2 and Fig. 4(a – g) present the results of each output flux in percentage terms. The model indicates that the increased N deposition increases the leaching of nitrate and ammonium from the catchment. The modelled results fit well with actual measurements for the PEFI catchment (Bicudo *et al.*, 2002), which endorses the value and appropriateness of the modelling. Nevertheless, the increased leaching does not account for all of the N deposition and so the catchment compensates for the increasing deposition in other ways. For example, the denitrification rates increase significantly, as might well be expected given the wet soils, warm temperatures and the sudden availability of excess N. Further, the nitrification processes also increase and so the excess ammonia is converted into nitrate. The analysis shows that nitrogen storage increases significantly (Table 3). This increase means that a larger pool of nitrogen is available in the catchment, increasing N concentration in the groundwater and this is available for future leaching.

Discussion and conclusions

This paper provides the first application of INCA to the South Americas and to the ecologically important area of the Atlantic Forest and it is hoped that the study will encourage new modelling initiatives in the region. The study raises important issues to be addressed; these include:

- The major need for high quality data in South American catchments to bring the level of understanding up to that known elsewhere (Neal and Heathwaite, 2005) and to take the science even further. Detailed information is generally sporadic and incomplete for the Americas. Process rates for nitrogen transformation are needed for model parameterisation to avoid using data from other parts of the world and calibration procedures rather than real measurements.




Fig. 4. Simulated parameter values variation for the different NO₃-N Wet deposition (a), NO₃-N Leaching (b), NO₃-N Uptake (c), NO₃-N denitrification (d), NH₄-N Total Load (e), NH₄-N Leaching (f), NH₄-N Nitrification (g) in kg.ha⁻¹.y

Table 3. Simulated, observed and N fluxes for increasing N deposition (kg N.ha⁻¹.yr⁻¹).

Parameter		0%	100%	200%	300%	400%	500%
NO ₃ -N deposition	simulated	9.09	18.0	27	36.0	45.0	54.0
NH ₄ -N deposition	simulated	8.07	18.15	27.2	36.3	45.38	54.45
NO ₃ -N leaching	simulated	6.5	8.26	9.83	11.41	12.99	14.57
	observed	5.22	-	-	-	-	-
NH ₄ -N leaching	simulated	3.85	5.19	6.39	7.60	8.80	10.01
	observed	2.38	-	-	-	-	-
NH ₄ -N nitrification	simulated	15.79 (pool)	23.69	30.8	37.91	45.02	52.15
NO ₃ -N denitrification	simulated	9.92	12.91	15.61	18.3	21.0	27.70
D Store	simulated	3.11	9.79	22.4	35.3	47.6	56.2

- Intensive catchment studies such as those in temperate regions of the world, in the UK at Plynlimon (Neal, 1997, 2002) and in the USA at Hubbard Brook (Likens and Bormann, 1995), need to be established and maintained in tropical environments to provide baseline

- data for environmental change studies.
- The McIntyre *et al.* (2005) sensitivity analysis using INCA-N showed the importance of investigating semi-distributed model uncertainties prior to model application. This work should be repeated for tropical

applications of INCA where process rates and interactions may differ from those in temperate regions.

Application of INCA facilitates analysis of the patterns of NO₃-N concentrations in discharge over the year and under varying hydrological conditions. It is a valuable tool for estimating NO₃-N concentrations and fluxes under diverse deposition backgrounds and hydrological regimes, as well as for water resources management. However, its usefulness needs to be tested in broader tropical regions with different soil uses and over a longer period of observations.

Acknowledgements

One author (Ranzini) thanks CNPq for a post-doctoral grant - N°150.350/03-1. The dataset were from FAPESP project N°99/05204-4. The authors thank J.B.A. dos Santos (Forestry Institute of São Paulo State) for his long-lasting help in field sampling and FAPESP Process No. 99/5204-4 for financial support on the hydrogeochemistry.

References

- Arcova, F.C.S., 1996. *Balanço hídrico, características do deflúvio e calibragem de duas microbacias hidrográficas na Serra do Mar, SP* [in Portuguese, with English abstract]. Msc Dissertation. Universidade de São Paulo. 155pp.
- Bastrup-Birk, A. and Gundersen, P., 2004. Water quality improvements from afforestation in an agricultural catchment in Denmark illustrated with the INCA model. *Hydrol. Earth Syst. Sci.*, **8**, 764–777.
- Bicudo, D.C., Forti, M.C., Carmo, C.F., Bourotte, C., Bicudo, C.E.M., Melfi, A. J. and Lucas, Y., 2002. A atmosfera, as águas superficiais e os reservatórios do PEFI. In: *Parque Estadual das Fontes do Ipiranga (PEFI): uma unidade de conservação ameaçada pela urbanização de São Paulo*, D.C. Bicudo, M.C. Forti and C.E.M. Bicudo (Eds). [in Portuguese] Ed. da Secretaria do Meio Ambiente do Estado de São Paulo, Cap 10:161–200.
- Camargo, A.P., 1971. Balanço hídrico no Estado de São Paulo [in Portuguese]. *Bol. Tecn. IAC*, **116**. 24pp.
- Cicco, V., 2004. *Análise de séries temporais hidrológicas em microbacia com cobertura vegetal natural de mata atlântica, Cunha – SP*. [in Portuguese, with English abstract]. MSc Dissertation. Universidade de São Paulo. 124pp.
- Emplasa (Empresa Paulista de Planejamento Metropolitano), 2005. *População Metropolitana Estimada em Tempo Real* [in Portuguese]. <http://www.emplasa.sp.gov.br>.
- Forti, M.C., Boulet, R., Melfi, A.J. and Neal, C., 2000. Hydrogeochemistry of a small catchment in north eastern Amazonia: a comparison between natural with deforested parts of the catchment (Serra do Navio, Amapá State, Brazil). *Water Air Soil Pollut.*, **118**, 263–279.
- Forti, M.C., Bicudo, D.C., Bourotte, C., Cicco, V. and Arcova, F.C.S., 2005. Rainfall and throughfall chemistry in the Atlantic Forest: a comparison between urban and natural areas (São Paulo State, Brazil). *Hydrol. Earth Syst. Sci.*, **9**, 570–585.
- Fujieda, F., Kudoh, T., Cicco, V. and Carvalho, J.L., 1997. Hydrological processes at two subtropical forest catchments: the Serra do Mar, São Paulo, Brazil. *J. Hydrol.*, **196**, 26–46.
- Fundação SOS Mata Atlântica, 2005. *Atlas dos remanescentes florestais da Mata Atlântica: período de 1995-2000*. Relatório Final. http://201.6.118.168/sos/download/relatorio_atlas-1995a2000.pdf. 1–43
- Furian, S.M. and Pfeifer, R.M., 1986. Levantamento de reconhecimento do meio físico do Núcleo Cunha, SP [in Portuguese]. *Bol. Tecn. IF*, **40**, 183–193.
- Jarvie, H.P., Wade, A.J., Butterfield, D., Whitehead, P.G., Tindall, C.I., Virtue, W.A., Dryburgh, W. and McGraw, A., 2002. Modelling nitrogen dynamics and distributions in the River Tweed, Scotland: an application of the INCA model. *Hydrol. Earth Syst. Sci.*, **6**, 433–453.
- Langusch, J.-J. and Matzner, E. N., 2002. Fluxes in two nitrogen saturated forested catchments in Germany: Dynamics and modelling with INCA. *Hydrol. Earth Syst. Sci.*, **6**, 383–394.
- Likens, G.E. and Borman, F.H., 1995. *Biogeochemistry of a Forested Ecosystem*. Springer, New York, 2nd Edn. 159pp.
- McIntyre, N., Jackson, B., Wade, A.J., Butterfield, D. and Wheeler, H.S., 2005. Sensitivity analysis of a catchment-scale nitrogen model. *J. Hydrol.*, **315**, 71–92.
- Nash, J.E. and Sutcliffe, J.V., 1970. River flow forecasting through conceptual models, 1. A discussion of principles. *J. Hydrol.*, **10**, 282–290.
- Neal, C., 1997. A view of water quality from the Plynlimon watershed. *Hydrol. Earth Syst. Sci.*, **1**, 743–753.
- Neal, C., 2002. Nutrient concentrations and fluxes for podzolic and gley soils at Plynlimon, mid-Wales: implications for modelling inorganic nitrogen and phosphorus in upland UK environments. *Hydrol. Earth Syst. Sci.*, **6**, 403–420.
- Neill, C., Piccolo, M.C., Cerri, C.C., Stender, P.A., Melillo, J.M. and Britto, M., 1997. Net nitrogen mineralization and net nitrification rates in soils following deforestation for pasture across south western Brazilian Amazon landscape. *Oecologia*, **110**, 243–252.
- Ranzini, M., 2002. *Modelagem hidrológica de uma microbacia florestada da serra do mar, SP, com o modelo TOPMODEL — simulação do comportamento hidrológico em função do corte raso* [in Portuguese, with English abstract]. Doctoral Thesis. Universidade de São Paulo. 116pp.
- Ranzini, M., Righetto, A.M., Lima, W.P., Guandique, M.E.G., Arcova, F.C.S. and Cicco, V., 2004. Processos hidrológicos de uma microbacia com Mata Atlântica, na região da Serra do Mar, SP [in Portuguese, with English abstract]. *Sci. forestalis*, **66**, 108–119.
- Wade, A.J., Durand, P., Beaujouan, V., Wessel, W.W., Raat, K.J., Whitehead, P.G., Butterfield, D., Rankinen, K. and Lepisto, A., 2002. A nitrogen model for European catchments: INCA, new model structure and equations. *Hydrol. Earth Syst. Sci.*, **6**, 559–582.
- Whitehead, P.G., Wilson, E.J. and Butterfield, D., 1998a. A semi-distributed integrated nitrogen model for multiple source assessment in catchments (INCA): part I — model structure and process equations. *Sci. Total Environ.*, **210/211**, 547–558.
- Whitehead, P.G., Wilson, E.J., Butterfield, D. and Seed, K., 1998b. A semi-distributed integrated flow and nitrogen model for multiple source assessment in catchments (INCA): part II — application to large river basins in south Wales and eastern England. *Sci. Total Environ.*, **210/211**, 559–583.
- Whitehead, P.G., Lapworth, D.J., Skeffington, R.A. and Wade, A., 2002. Excess nitrogen leaching and C/N decline in the Tillingbourne catchment, southern England: INCA process modelling for current and historic time series. *Hydrol. Earth Syst. Sci.*, **6**, 455–466.